

Existence of Chaos in Evolution Equations

Yanguang (Charles) Li *

School of Mathematics

Institute for Advanced Study

Princeton, NJ 08540

February 1, 2008

Abstract

For a general evolution equation with a Silnikov homoclinic orbit, Smale horseshoes are constructed with the tools of [1] and in the same way as in [1]. The linear part of the evolution equation has a finite number of unstable modes. For evolution equations with infinitely many linearly unstable modes, the problem is still open.

MSC: 35, 37.

1 Introduction

In recent years, the existence of chaos in partial differential equations has been established [2], [3], [1]. These studies set up a scheme for attacking problems on Chaos in PDEs. The type of chaos studied in these works is the so-called homoclinic chaos generated in a neighborhood of a homoclinic orbit. Two types of homoclinic orbits have been studied. One type is the so-called transversal homoclinic orbit [4]. The other type is the so-called Silnikov homoclinic orbit, which is non-transversal [2], [3], [1]. For lower dimensional and

*This work is supported by a Guggenheim Fellowship, and an AMS Centennial Fellowship. Current Address: Department of Mathematics, University of Missouri, Columbia, MO 65211. Email: cli@math.missouri.edu

general finite dimensional systems, Silnikov studied the symbolic dynamics structures in the neighborhoods of such homoclinic orbits [5], [6], [7], [8]. In [9] and [1], we have developed a different construction of Smale horseshoes in the neighborhood of a Silnikov homoclinic orbit. The advantages of our constructions have been fully addressed in [9], [1]. In this note, we generalize the construction in [1] to more general evolution equations with finitely many linearly unstable modes. For evolution equations with infinitely many linearly unstable modes, the problem is still open.

2 The Set-Up

Consider the evolution equation

$$(1) \quad \partial_t u = \mathcal{L}u + \mathcal{N}(u) ,$$

where $\mathcal{L}$ is a linear operator which is constant in time, and $\mathcal{N}$ is the nonlinear term.

Following are the assumptions for the setup:

(A1) $u = 0$ is a saddle, the linear operator $\mathcal{L}$ has only point spectrum as follows,

$$\sigma(\mathcal{L}) = \{ -\alpha \pm i\beta, \gamma, \lambda_j^\pm, j \in S^\pm \subset Z^+ \} ,$$

where $\alpha > 0, \beta > 0, \gamma > 0, \operatorname{Re} \{ \lambda_j^\pm \} \geq 0$; the number of elements in S^+ is finite, denoted by N , $\alpha < \gamma$, and

$$\alpha < \lambda^- = \inf_{j \in S^-} \{ -\operatorname{Re} \{ \lambda_j^- \} \} , \quad \gamma < \lambda^+ = \inf_{j \in S^+} \{ \operatorname{Re} \{ \lambda_j^+ \} \} .$$

(A2) The evolution equation (1) is globally well posed in a Hilbert space $\mathcal{H}$, that is, there exists a unique solution to the Cauchy problem of (1), $u(t, u_0) \in C^0[(-\infty, \infty), \mathcal{H}]$, $u(0, u_0) = u_0$. Moreover, we assume the regularity condition on initial data that for any $t \in (-\infty, \infty)$, the evolution operator $F(u_0)^t = u(t, u_0)$ is C^n in u_0 for some $n \geq 2$.

(A3) With respect to the saddle $u = 0$, the evolution operator $F^t(u)$ admits a C^2 smooth linearization, i.e. there exists a C^2 diffeomorphism $\mathcal{R}: \mathcal{H} \rightarrow \mathcal{H}$, such that in terms

of the new variable $\tilde{u} = \mathcal{R}u$, the evolution equation (1) is transformed into the linear form

$$\partial_t \tilde{u} = \mathcal{L} \tilde{u}$$

in a neighborhood of $u = 0$. The conjugated evolution operator

$$\tilde{F}^t = \mathcal{R} F^t \mathcal{R}^{-1}$$

is still C^0 in time t and C^2 in $\tilde{u}$. In fact, we assume that the evolution equation (1) takes the normal form:

$$\begin{cases} \dot{x} = -\alpha x - \beta y + G_x(x, y, z, v^\pm), \\ \dot{y} = \beta x - \alpha y + G_y(x, y, z, v^\pm), \\ \dot{z} = \gamma z + G_z(x, y, z, v^\pm), \\ \partial_t v^\pm = L^\pm v^\pm + G_{v^\pm}(x, y, z, v^\pm), \end{cases}$$

where $G = 0$ in a neighborhood Ω of 0, $v^+ = (v_1^+, \dots, v_N^+)', N$ is the number of elements in S^+ ; x, y, z , and $v_j^\pm$'s are real variables, and

$$\|e^{L^+ t}\| \leq c^+ e^{\lambda^+ t}, \quad \text{as } t \rightarrow -\infty, \quad \|e^{L^- t}\| \leq c^- e^{-\lambda^- t}, \quad \text{as } t \rightarrow +\infty.$$

For references on such linearization results, see for example [10] etc.

- (A4) There exists a Silnikov homoclinic orbit $h(t)$ asymptotic to 0. As $t \rightarrow +\infty$, h is tangent to the (x, y) -plane at 0, and as $t \rightarrow -\infty$, h is tangent to the z -axis (without loss of generality, positive z -axis) at 0. The stable and unstable manifolds of 0 are C^2 smooth, and

$$\dim \{\mathcal{T}_v W^u \cap \mathcal{T}_v W^s\} = 1,$$

where $v \in h(t)$, $\mathcal{T}_v W^u$ is the tangent space of the unstable manifold of 0 at the point v on the homoclinic orbit $h(t)$, similarly for $\mathcal{T}_v W^s$.

Remark. Proving the existence of Silnikov homoclinic orbits in partial differential equations is a rather nontrivial question. So far, this has been done for perturbed nonlinear Schrödinger equation [3] [11], perturbed vector nonlinear Schrödinger equation [12], and perturbed discrete nonlinear Schrödinger equation [13]. The above and later assumptions have been either

verified or discussed for these equations in [1] [14] [9]. The perturbed Davey-Stewartson II equation has been studied along this direction [15] [16]. Unfortunately, existence of Silnikov homoclinic orbits has not been proved due to some technical difficulty [16]. I would like to comment on equations that have the potential of being casted into the above setup: 1. perturbations of the modified KdV equation

$$u_t + 6u^2u_x + u_{xxx} = 0 ,$$

2. perturbations of the derivative nonlinear Schrödinger equation [17] [18] [19] [20]

$$iu_t = u_{xx} + i\alpha(|u|^2u)_x + 2|u|^2u , \quad \alpha > 0 ,$$

and 3. perturbations of the derivative nonlinear Schrödinger equation

$$iu_t = u_{xx} - i\alpha u^2 \bar{u}_x + 2|u|^2u + 2\alpha^2|u|^4u .$$

All of the above three equations are integrable systems.

3 The Construction of Smale Horseshoes

3.1 Definitions

Definition. The Poincaré section Σ_0 is defined by the constraints:

$$y = 0, \quad \eta \exp\{-2\pi\alpha/\beta\} < x < \eta ,$$

$$0 < z < \eta, \quad \|v^\pm\| < \eta ;$$

where η is a sufficiently small constant so that Σ_0 is included in the neighborhood Ω of 0 where the dynamics is given by the linear system.

Definition. The auxiliary section Σ_0^+ is defined by the constraints:

$$y = 0, \quad \eta \exp\{-2\pi\alpha/\beta\} < x < \eta ,$$

$$-\eta < z < \eta, \quad \|v^\pm\| < \eta .$$

The homoclinic orbit h intersects the $(z = 0)$ -boundary of $\sum_0$ at $w^{(+)}$ with coordinates denoted by

$$x = x_*, \ y = 0, \ z = 0, \ v^+ = 0, \ v^- = v_*^- .$$

There exists $T > 0$ such that the point $w^{(-)} = F^{-T}(w^{(+)})$ on h (where F^t is the evolution operator) has the z coordinate equal to η . Denote the coordinates of $w^{(-)}$ by

$$x = y = v^- = 0, \ z = \eta, \ v^+ = v_*^+ .$$

Definition. The Poincaré section $\sum_1$ is defined as:

$$\sum_1 = \left(F^{-T} \circ \sum_0^+ \right) \cap \Omega .$$

Definition. The map P_0^1 from $\sum_0$ to $\sum_1$ is defined as:

$$\begin{aligned} P_0^1: \ U_0 \subset \sum_0 &\mapsto \sum_1 , \\ \forall w \in U_0, \ P_0^1(w) &= F^{t_*}(w) \in \sum_1 , \end{aligned}$$

where $t_* = t_*(w) > 0$ is the smallest time t such that $F^t(w) \in \sum_1$.

The map from $\sum_1$ to $\overline{\sum_0} (= \sum_0 \cup \partial \sum_0)$ is defined as:

$$\begin{aligned} P_1^0: \ U_1 \subset \sum_1 &\mapsto \overline{\sum_0} , \\ \forall w \in \sum_1, \ P_1^0(w) &= F^T(w) \in \overline{\sum_0} . \end{aligned}$$

The Poincaré map P from $\sum_0$ to itself is defined as:

$$\begin{aligned} P: \ U \subset \sum_0 &\mapsto \sum_0 , \\ P &= P_1^0 \circ P_0^1 . \end{aligned}$$

3.2 Fixed Points of the Poincaré Map P

On the Poincaré section $\sum_0$, we center the origin of the coordinate frame at $w^{(+)}$, and denote the new coordinates by $(x^{(0)}, z^{(0)}, v^{(\pm,0)})$. On the Poincaré section $\sum_1$, we center the origin of the coordinate frame at $w^{(-)}$, and denote the new coordinates by $(x^{(1)}, y^{(1)}, z^{(1)}, v^{(\pm,1)})$, which satisfy the constraint equation

$$(2) \quad f^{(y)} \left(x^{(1)}, y^{(1)}, z^{(1)} + \eta, v^{(+,1)} + v_*^+, v^{(-,1)} \right) = 0$$

where for any $w \in \mathcal{H}$,

$$F^T(w) = \left(f^{(x)}(w), f^{(y)}(w), f^{(z)}(w), f^{(v^+)}(w), f^{(v^-)}(w) \right) .$$

Denote the vector v^+ in component form, $v^+ = (v_1^+, v_2^+, \dots, v_N^+)'$. Then we have the lemma,

Lemma 1. $\frac{\partial f^{(y)}}{\partial z}(w^{(-)})$ and $\frac{\partial f^{(y)}}{\partial v_j^+}(w^{(-)})$ ($j = 1, \dots, N$) cannot be zero simultaneously.

Proof. Assume that they are zero simultaneously, then

$$\tau \cdot \nabla f^{(y)}(w^{(-)}) = 0 ,$$

where τ is the tangent vector of h at $w^{(-)}$ and “ ∇ ” denotes gradient. Then this implies that h is tangent to $\sum_0$ at $w^{(-)}$. This contradiction proves the lemma. $\square$

Let ξ be one of the coordinates $\{z, v_j^+ (j = 1, \dots, N)\}$, such that $\frac{\partial f^{(y)}}{\partial \xi}(w^{(-)}) \neq 0$, and denote by $v_+^{(1)}$ the vector $(z^{(1)}, v_j^{(+,1)} (j = 1, \dots, N))' \setminus \{\xi\}$, i.e. with components consisting of $\{z^{(1)}, v_j^{(+,1)} (j = 1, \dots, N)\}$ without ξ .

Lemma 2. In a neighborhood of $w^{(-)}$, the Poincaré section $\sum_1$ can be represented as a C^2 function

$$\xi = \xi \left(x^{(1)}, y^{(1)}, v^{(-,1)}, v_+^{(1)} \right) .$$

Proof. Applying the implicit function theorem to (2). $\square$

The map P_0^1 has the representation

$$\begin{cases} x^{(1)} = e^{-\alpha t_*} (x^{(0)} + x_*) \cos \beta t_*, \\ y^{(1)} = e^{-\alpha t_*} (x^{(0)} + x_*) \sin \beta t_*, \\ z^{(1)} + \eta = z^{(0)} e^{\gamma t_*}, \\ v^{(+,1)} + v_*^+ = e^{L^+ t_*} v^{(+,0)}, \\ v^{(-,1)} = e^{L^- t_*} (v^{(-,0)} + v_*^-) . \end{cases}$$

The map P_1^0 can be approximated by its linearization at $w^{(-)}$,

$$\begin{pmatrix} x^{(0)} \\ z^{(0)} \\ v^{(+,0)} \\ v^{(-,0)} \end{pmatrix} = A \begin{pmatrix} x^{(1)} \\ y^{(1)} \\ z^{(1)} \\ v^{(+,1)} \\ v^{(-,1)} \end{pmatrix},$$

where

$$A = \begin{pmatrix} \frac{\partial f^{(x)}}{\partial x} & \frac{\partial f^{(x)}}{\partial y} & \frac{\partial f^{(x)}}{\partial z} & \frac{\partial f^{(x)}}{\partial v^+} & \frac{\partial f^{(x)}}{\partial v^-} \\ \frac{\partial f^{(z)}}{\partial x} & \frac{\partial f^{(z)}}{\partial y} & \frac{\partial f^{(z)}}{\partial z} & \frac{\partial f^{(z)}}{\partial v^+} & \frac{\partial f^{(z)}}{\partial v^-} \\ \frac{\partial f^{(v^+)}}{\partial x} & \frac{\partial f^{(v^+)}}{\partial y} & \frac{\partial f^{(v^+)}}{\partial z} & \frac{\partial f^{(v^+)}}{\partial v^+} & \frac{\partial f^{(v^+)}}{\partial v^-} \\ \frac{\partial f^{(v^-)}}{\partial x} & \frac{\partial f^{(v^-)}}{\partial y} & \frac{\partial f^{(v^-)}}{\partial z} & \frac{\partial f^{(v^-)}}{\partial v^+} & \frac{\partial f^{(v^-)}}{\partial v^-} \end{pmatrix}_{(w^{(-)})}.$$

The constraint equation (2) can be approximated by its linearization at $w^{(-)}$,

$$B \begin{pmatrix} x^{(1)} \\ y^{(1)} \\ z^{(1)} \\ v^{(+,1)} \\ v^{(-,1)} \end{pmatrix} = 0,$$

where

$$B = \left(\frac{\partial f^{(y)}}{\partial x} \frac{\partial f^{(y)}}{\partial y} \frac{\partial f^{(y)}}{\partial z} \frac{\partial f^{(y)}}{\partial v^+} \frac{\partial f^{(y)}}{\partial v^-} \right)_{(w^{(-)})}.$$

With the above preparations, we can write the equations of the fixed points of P as follows in terms of the Silnikov coordinates $\{t_*, x^{(0)}, v^{(-,0)}, z^{(1)}, v^{(+,1)}\}$

$$(3) \quad \begin{pmatrix} x^{(0)} \\ (z^{(1)} + \eta)e^{-\gamma t_*} \\ e^{-L^+ t_*}(v^{(+,1)} + v_*^+) \\ v^{(-,0)} \end{pmatrix} = A \begin{pmatrix} e^{-\alpha t_*}(x^{(0)} + x_*) \cos \beta t_* \\ e^{-\alpha t_*}(x^{(0)} + x_*) \sin \beta t_* \\ z^{(1)} \\ v^{(+,1)} \\ e^{L^- t_*}(v^{(-,0)} + v_*^-) \end{pmatrix} + \mathcal{R},$$

where $\mathcal{R} = \mathcal{O}(e^{-\nu t_*})$ as $t_* \rightarrow +\infty$ for some $\nu > 0$. By rescaling the coordinates as follows,

$$\{t_*, \hat{x}^{(0)} = x^{(0)} e^{\alpha t_*}, \hat{v}^{(-,0)} = v^{(-,0)} e^{\alpha t_*}, \hat{z}^{(1)} = z^{(1)} e^{\alpha t_*}, \hat{v}^{(+,1)} = v^{(+,1)} e^{\alpha t_*}\} ,$$

we can rewrite equation (3) in the form

$$(4) \quad \begin{pmatrix} \hat{x}^{(0)} \\ 0 \\ 0 \\ \hat{v}^{(-,0)} \end{pmatrix} = A \begin{pmatrix} x_* \cos \beta t_* \\ x_* \sin \beta t_* \\ \hat{z}^{(1)} \\ \hat{v}^{(+,1)} \\ 0 \end{pmatrix} + \mathcal{R}_1 ,$$

where $\mathcal{R}_1 = \mathcal{O}(e^{-\nu_1 t_*})$ as $t_* \rightarrow +\infty$ for some $\nu_1 > 0$, and the constraint equation (2) takes the form

$$(5) \quad B \begin{pmatrix} x_* \cos \beta t_* \\ x_* \sin \beta t_* \\ \hat{z}^{(1)} \\ \hat{v}^{(+,1)} \\ 0 \end{pmatrix} + \mathcal{R}_2 = 0 ,$$

where $\mathcal{R}_2 = \mathcal{O}(e^{-\nu_2 t_*})$ as $t_* \rightarrow +\infty$ for some $\nu_2 > 0$. Solving the leading order term of (4) for $(\hat{z}^{(1)}, \hat{v}^{(+,1)})$, we have

$$(6) \quad C \begin{pmatrix} \hat{z}^{(1)} \\ \hat{v}^{(+,1)} \end{pmatrix} = -D \begin{pmatrix} x_* \cos \beta t_* \\ x_* \sin \beta t_* \end{pmatrix} ,$$

where

$$C = \begin{pmatrix} \frac{\partial f(z)}{\partial z} & \frac{\partial f(z)}{\partial v^+} \\ \frac{\partial f(v^+)}{\partial z} & \frac{\partial f(v^+)}{\partial v^+} \end{pmatrix}_{(w^{(-)})} , \quad D = \begin{pmatrix} \frac{\partial f(z)}{\partial x} & \frac{\partial f(z)}{\partial y} \\ \frac{\partial f(v^+)}{\partial x} & \frac{\partial f(v^+)}{\partial y} \end{pmatrix}_{(w^{(-)})} .$$

Lemma 3. *The matrix C is invertible.*

Proof. Assume that C is non-invertible; then there exists a nonzero vector $\begin{pmatrix} \hat{z}^{(1)} \\ \hat{v}^{(+,1)} \end{pmatrix}$, such that

$$C \begin{pmatrix} \hat{z}^{(1)} \\ \hat{v}^{(+,1)} \end{pmatrix} = 0 ;$$

thus

$$w = A \begin{pmatrix} 0 \\ 0 \\ \hat{z}^{(1)} \\ \hat{v}^{(+,1)} \\ 0 \end{pmatrix} \in \mathcal{T}_{w^{(+)}} W^s \cap \mathcal{T}_{w^{(+)}} W^u$$

where $\mathcal{T}_{w^{(+)}} W^s$ is the tangent space of W^s at $w^{(+)}$, similarly for $\mathcal{T}_{w^{(+)}} W^u$. Since A is invertible, $w \neq 0$. We also know that

$$\mathcal{T}_{w^{(+)}} h \in \mathcal{T}_{w^{(+)}} W^s \cap \mathcal{T}_{w^{(+)}} W^u .$$

Since $\mathcal{T}_{w^{(+)}} h$ is transversal to $\sum_0$ and w lies in $\sum_0$, we have

$$\dim \{ \mathcal{T}_{w^{(+)}} W^u \cap \mathcal{T}_{w^{(+)}} W^s \} = 2 ,$$

which contradicts with Assumption (A4). This completes the proof. $\square$

Solving (6), we have

$$(7) \quad \begin{pmatrix} \hat{z}^{(1)} \\ \hat{v}^{(+,1)} \end{pmatrix} = -C^{-1} D \begin{pmatrix} x_* \cos \beta t_* \\ x_* \sin \beta t_* \end{pmatrix} .$$

Then solving the constraint equation (5), to the leading order, we have

$$B \begin{pmatrix} x_* \cos \beta t_* \\ x_* \sin \beta t_* \\ -C^{-1} D \begin{pmatrix} x_* \cos \beta t_* \\ x_* \sin \beta t_* \end{pmatrix} \\ 0 \end{pmatrix} = 0 ,$$

which can be rewritten as

$$(8) \quad \Delta_1 \cos \beta t_* + \Delta_2 \sin \beta t_* = 0 .$$

If we assume condition

(A5) Δ_1 and Δ_2 do not vanish simultaneously,

then (8) has a sequence of solutions:

$$(9) \quad t_*^{(\ell)} = \frac{1}{\beta} [\ell\pi - \varphi], \quad \ell \in Z^+,$$

where

$$\varphi = \arctan \{ \Delta_1 / \Delta_2 \} .$$

Then substituting (7) and (9) into the leading order terms of (4), we can solve for $\begin{pmatrix} \hat{x}^{(0)} \\ \hat{v}^{(-,0)} \end{pmatrix}$ to the leading order. Finally, applying the implicit function theorem, we have the fixed point theorem.

Theorem 1 (Fixed Point Theorem). *Under the assumption (A5), there exists an integer $\ell_0 > 0$, such that there exists a sequence of solutions to the equations (4) and (5) labeled by ℓ ($\ell \geq \ell_0$):*

$$\begin{aligned} t_* &= T^{(\ell)}, \quad \hat{x}^{(0)} = x^{(\ell)}, \quad \hat{v}^{(-,0)} = v_{(-,0)}^{(\ell)} \\ \hat{z}^{(1)} &= z^{(\ell)}, \quad \hat{v}^{(+,1)} = v_{(+,1)}^{(\ell)} ; \end{aligned}$$

where, as $t \rightarrow +\infty$,

$$T^{(\ell)} = \frac{1}{\beta} [\ell\pi - \varphi] + o(1) .$$

For a complete proof of this theorem, see [1].

3.3 Smale Horseshoes

Definition. For sufficiently large number ℓ , we define the slab S_ℓ in Σ_0 as follows:

$$\begin{aligned} S_\ell &= \left\{ w \in \Sigma_0 \mid \eta \exp \left\{ -\gamma \left(T^{(2(\ell+1))} - \frac{\pi}{2\beta} \right) \right\} \leq z^{(0)}(w) \right. \\ &\quad \leq \eta \exp \left\{ -\gamma \left(T^{(2\ell)} - \frac{\pi}{2\beta} \right) \right\} , \\ &\quad |x^{(0)}(w)| \leq \eta \exp \left\{ -\frac{1}{2} \alpha T^{(2\ell)} \right\} , \\ &\quad \|v^{(-,1)}(P_0^1(w))\| \leq \eta \exp \left\{ -\frac{1}{2} \alpha T^{(2\ell)} \right\} , \\ &\quad \left. \|v^{(+,1)}(P_0^1(w))\| \leq \eta \exp \left\{ -\frac{1}{2} \alpha T^{(2\ell)} \right\} \right\} , \end{aligned}$$

so that it contains two fixed points of P denoted by p_ℓ^+ and p_ℓ^- .




Figure 1: The product representation of S_ℓ .

We choose a basis on the tangent space $\mathcal{T}_{w^{(-)}} \sum_1$ represented in the coordinates

$$(x^{(1)}, y^{(1)}, v^{(-,1)}, \xi, v_+^{(1)})$$

as follows:

$$\begin{aligned} E_{x^{(1)}} &= (1, 0, 0, \nabla \xi(w^{(-)}) \circ (1, 0, 0, 0), 0) , \\ E_{y^{(1)}} &= (0, 1, 0, \nabla \xi(w^{(-)}) \circ (0, 1, 0, 0), 0) , \\ \mathbf{E}_{v^{(-,1)}} &= (0, 0, \mathbb{1}, \nabla \xi(w^{(-)}) \circ (0, 0, \mathbb{1}, 0), 0) , \\ \mathbf{E}_{v_+^{(1)}} &= (0, 0, 0, \nabla \xi(w^{(-)}) \circ (0, 0, 0, \mathbb{1}), \mathbb{1}) ; \end{aligned}$$

where $\mathbb{1}$ represents a basis for the corresponding components. Denote by $\{e_{x^{(0)}}, e_{z^{(0)}}, \mathbf{e}_{v^{(+,0)}}, \mathbf{e}_{v^{(-,0)}}\}$ the unit vectors along $(x^{(0)}, z^{(0)}, v^{(+,0)}, v^{(-,0)})$ -directions in $\sum_0$. In this coordinate frame, S_ℓ has the product representation as shown in Fig.1. Under the linear map A (the linearization of P_1^0 at $w^{(-)}$), the coordinate frame $\{E_{x^{(1)}}, E_{y^{(1)}}, \mathbf{E}_{v^{(-,1)}}, \mathbf{E}_{v_+^{(1)}}\}$ is mapped into a coordinate frame $\{\mathcal{E}_{x^{(1)}}, \mathcal{E}_{y^{(1)}}, \mathbf{\mathcal{E}}_{v^{(-,1)}}, \mathbf{\mathcal{E}}_{v_+^{(1)}}\}$ on $\sum_0$ with origin at $w^{(+)}$. In this coordinate frame, $\nabla P_1^0(w^{(-)}) \circ \nabla P_1^0(w^{(+)}) \circ S_\ell$ has the representation as shown in Fig.2 on $\sum_0$. We introduce a system of curvilinear coordinates (ξ_u, ξ_s) on the $(\mathcal{E}_{x^{(1)}}, \mathcal{E}_{y^{(1)}})$ -plane such that

$$\{\xi_u = 0\}, \{\xi_u = c_u \text{ (a constant)}\}, \{\xi_s = 0\}, \{\xi_s = c_s \text{ (a constant)}\}$$

correspond to the boundaries 3, 4, 1, 2 of the annulus on the $(\mathcal{E}_{x^{(1)}}, \mathcal{E}_{y^{(1)}})$ -plane. Let $\mathcal{E}_\xi^+ = \mathcal{T}_{p_\ell^+} \xi_u$ be the tangent vector to the ξ_u coordinate at p_ℓ^+ . We make the following assumption:

$$(A6) \quad \text{Span} \{e_{x^{(0)}}, \mathbf{e}_{v^{(-,0)}}, \mathcal{E}_\xi^+, \mathbf{\mathcal{E}}_{v_+^{(1)}}\} = \sum_0 .$$




Figure 2: A geometric representation of $\nabla P_1^0(w^{(-)}) \circ \nabla P_0^1(w^{(+)}) \circ S_\ell$.

Under the assumptions (A1) – (A6), we can verify the Conley-Moser conditions [21] in the same way as in [1], which lead to the construction of Smale horseshoes. Let $\mathcal{W}$ be a set which consists of elements of the doubly infinite sequence form

$$a = (\cdots a_{-2}a_{-1}a_0, a_1a_2\cdots),$$

where $a_k \in \{0, 1\}$, $k \in Z$. We introduce a topology in $\mathcal{W}$ by taking as neighborhood basis of

$$a^* = (\cdots a_{-2}^* a_{-1}^* a_0^*, a_1^* a_2^* \cdots),$$

the set

$$W_j = \{a \in \mathcal{W} \mid a_k = a_k^* (|k| < j)\}$$

for $j = 1, 2, \dots$. This makes $\mathcal{W}$ a topological space. The shift automorphism χ is defined on $\mathcal{W}$ by

$$\chi \quad : \quad \mathcal{W} \longmapsto \mathcal{W} \, ,$$

$$\forall a \in \mathcal{W}, \chi(a) = b, \quad \text{where} \quad b_k = a_{k+1} .$$

The shift automorphism χ exhibits sensitive dependence on initial conditions, which is the *hallmark* of *chaos*.

Theorem 2 (Smale Horseshoe Theorem). *Under the assumptions (A1) – (A6) for the evolution equation (1), for all sufficiently large integers ℓ , there exists a sequence of compact Cantor subsets Λ_ℓ of S_ℓ , Λ_ℓ consists of points and is invariant under P . P restricted to Λ_ℓ is*

topologically conjugate to the shift automorphism χ on two symbols 0 and 1. That is, there exists a homeomorphism

$$\phi_\ell: \mathcal{W} \mapsto \Lambda_\ell ,$$

such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{W} & \xrightarrow{\phi_\ell} & \Lambda_\ell \\ \chi \downarrow & & \downarrow P \\ \mathcal{W} & \xrightarrow{\phi_\ell} & \Lambda_\ell . \end{array}$$

Proof. With preliminaries given above, the proof follows in the same way as in [1]. $\square$

4 Conclusion and Discussion

In this note, we have generalized the construction of Smale horseshoes in [1] to a general evolution equation, which indicates that our techniques [9], [1] on constructing Smale horseshoes in a neighborhood of a Silnikov homoclinic orbit has a much wider application. On the other hand, so far we can only handle evolution equations with finitely many linearly unstable modes. For evolution equations with infinitely many linearly unstable modes, we cannot invert certain linear operators in establishing the existence of fixed points of the Poincaré map. Nevertheless, this note furnishes an initiation for studying Silnikov homoclinic orbits for general evolution equations, thereby proving the existence of chaos for general evolution equations.

References

- [1] Y. Li, Smale horseshoes and symbolic dynamics in perturbed nonlinear Schrödinger equations, *Journal of Nonlinear Sciences* **9**, no.4 (1999), 363.
- [2] Y. Li and D. W. McLaughlin, Morse and Melnikov functions for NLS pdes, *Comm. Math. Phys.* **162** (1994), 175.
- [3] Y. Li et al., Persistent homoclinic orbits for a perturbed nonlinear Schrödinger equation, *Comm. Pure Appl. Math.* **XLIX** (1996), 1175.

- [4] Y. Li, Chaos and shadowing lemma for autonomous systems of infinite dimensions, *Submitted, available at: <http://www.math.missouri.edu/~cli>*, (2002).
- [5] L. P. Silnikov, A case of the existence of a countable number of periodic motions, *Soviet Math. Doklady* **6** (1965), 163.
- [6] L. P. Silnikov, The existence of a denumerable set of periodic motions in four-dimensional space in an extended neighborhood of a saddle-focus, *Soviet Math. Doklady* **6** (1965), 163.
- [7] L. P. Silnikov, On a Poincare-Birkoff problem, *Math. USSR Sb.* **3** (1967).
- [8] L. P. Silnikov, A contribution to the problem of the structure of an extended neighborhood of a rough equilibrium state of saddle-focus type, *Math. USSR Sb.* **10** (1970).
- [9] Y. Li and S. Wiggins, Homoclinic orbits and chaos in discretized perturbed NLS system, part II. symbolic dynamics, *Journal of Nonlinear Sciences* **7** (1997), 315.
- [10] N. V. Nikolenko, The method of Poincare normal forms in problems of integrability of equations of evolution type, *Russian Math. Surveys* **41:5** (1986), 63.
- [11] Y. Li, Persistent homoclinic orbits for nonlinear Schrödinger equation under singular perturbation, *Submitted* (2002).
- [12] Y. Li, Singularly perturbed vector and scalar nonlinear Schrödinger equations with persistent homoclinic orbits *Studies in Applied Mathematics* **current issue** (2002).
- [13] Y. Li and D. W. McLaughlin, Homoclinic orbits and chaos in discretized perturbed NLS system, part I. homoclinic orbits, *Journal of Nonlinear Sciences* **7** (1997), 211.
- [14] Y. Li, Existence of chaos for a singularly perturbed NLS equation, *Submitted* (2002).
- [15] Y. Li, Bäcklund-Darboux transformations and Melnikov analysis for Davey-Stewartson II equations, *Journal of Nonlinear Sciences* **10, no.1** (2000), 103.
- [16] Y. Li, Melnikov analysis for singularly perturbed DSII equation, *Submitted* (2002).

- [17] V. S. Shchesnovich and E. V. Doktorov, Perturbation theory for the modified nonlinear Schroedinger solitons, *Phys. D* **129**, no.1-2 (1999), 115.
- [18] V. S. Gerdjikov and E. V. Doktorov and J. Yang, Adiabatic interaction of N ultrashort solitons: universality of the complex Toda chain model, *Phys. Rev. E* **64**, no.5, part 2 (2001).
- [19] K. Mio et al., Modified nonlinear Schroedinger equation for Alfvén waves propagating along the magnetic field in cold plasmas, *J. Phys. Soc. Japan* **41**, no.1 (1976), 265.
- [20] M. Wadati et al., A generalization of inverse scattering method, *J. Phys. Soc. Japan* **46**, no.6 (1979), 1965.
- [21] J. Moser, Stable and Random Motions in Dynamical Systems, *Annals of Mathematics Studies, Princeton University Press* **77** (1973).